

University of Groningen

Challenges in the use of preventive cardiovascular medications in Indonesia and the Netherlands

Irawati, Sylvi

DOI:
[10.33612/diss.146680004](https://doi.org/10.33612/diss.146680004)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Irawati, S. (2020). *Challenges in the use of preventive cardiovascular medications in Indonesia and the Netherlands*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.146680004>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ADDENDUM

SUMMARY

NEDERLANDSE

SAMENVATTING

ACKNOWLEDGMENT

ABOUT THE AUTHOR

PHD. PORTOFOLIO

SUMMARY

Cardiovascular disease (CVD) is a leading cause of morbidity and mortality worldwide. More than 75% of CVD-related deaths, mainly due to coronary heart disease (CHD) and stroke, take place in the low and middle-income countries (LMICs), including Indonesia. While CVD is no longer the main cause of hospitalizations or death in the Netherlands, one of the high-income countries (HICs), it is still the primary cause of death in Indonesia. To reduce the burden of CVD, the World Health Organization (WHO) and international cardiovascular (CV)-related organizations issue clinical guidelines that recommend, amongst other measures, the use of preventive CV medications based on an individual CV risk assessment. There are several classes of preventive CV medications recommended with evidence of effectiveness in CVD prevention and control. These medications are antithrombotic agents (aspirin, P₁₂Y₁₂ inhibitors, anticoagulants), antihypertensive agents (e.g., beta blockers, angiotensin-converting enzyme inhibitors [ACEIs], angiotensin receptor blockers [ARBs]) and lipid lower agents, e.g., statins. To optimize the use of these preventive medications in populations at risk remains a challenge. These challenges are different in Indonesia as an LMIC and the Netherlands as an HIC. Studies on the effectiveness of these medications in an Indonesian population at risk of CVD are very limited. It has been reported that CVD-related guidelines and CV risk stratification are underutilized. Within the new National Health Insurance System (NHIS, namely *Jaminan Kesehatan Nasional-Kartu Indonesia Sehat* [JKN-KIS]) in Indonesia, the primary healthcare sector plays an important role to influence the use of preventive CV medications. Meanwhile, systematic and programmatic CVD prevention in the Netherlands has already been introduced since 2005. The role of the Dutch primary healthcare sector and health insurance companies in CVD prevention is prominent. The clinical guidelines to manage multi CV risk factors have already been developed since the mid-2000s. The differences in the stage in CVD prevention between both countries may result in distinct challenges related to the use of preventive CV medications in Indonesia and the Netherlands. This thesis aims to identify current challenges in the use of guideline-recommended CV preventive medications in Indonesia and the Netherlands. CHD is the type of CVD that we focus on. The thesis is divided into two parts: **Part 1 (Chapter 2-5)** investigates the long-term CVD burden in Asians and the challenges in using preventive CV medications in Indonesia, and **Part 2 (Chapter 6 and 7)** studies the challenges in using specific preventive CV medications in the Netherlands.

In **Chapter 2**, a systematic review and meta-analysis was conducted to estimate the long-term (> 10 year) risk of CVD and its risk factors in Asian populations. Most studies identified were from the East-Asian region, none was from Indonesia. The long-term risk of fatal CVD in the Asian populations was 6.35% during a 20.00 years of mean follow-up. The long-term risk of fatal stroke was higher than of fatal CHD. Different risk factors between CHD and stroke were identified. Men, older age and current smokers were risk factors for fatal CVD identified from the meta-analysis. In contrast to previous studies in Western populations, we concluded that different approaches might be necessary for CVD prevention and control in Asians due to the different burden and risk factors compared to Westerners.

In **Chapter 3**, a cohort study aimed to estimate the effectiveness of guideline-recommended preventive CV medications on in-hospital mortality in patients with ST-elevation myocardial infarction (STEMI), an acute manifestation of CHD, was conducted using the Jakarta Acute Coronary Syndrome (JAC) Registry. We observed the use of preventive medications in 42% patients with STEMI not treated with acute primary percutaneous coronary intervention (pPCI) therapy, the majority of whom were late presenters at the hospital. A third of these patients did not receive guideline-recommended preventive medications at hospital admission. After potential confounders were taken into account in the statistical analysis, it was estimated that the use of preventive CV medications in this population reduced the odds of in-hospital death by more than 50%.

In **Chapter 4**, using a cohort study from the same registry, we assessed predictors for suboptimal use of guideline-recommended preventive CV medications in patients with STEMI. We found the predictors were the presence of non-anterior MI; age > 65 years; not being treated with acute reperfusion therapy; having a family history of CAD; and having a thrombolysis in myocardial infarction (TIMI) score ≥ 4 . Not being treated with reperfusion was identified as the common predictor for not receiving most drug classes separately.

From both **Chapter 4** and **5**, it seemed there was a presence of treatment-risk paradox, a situation where patients with higher risk tended to receive less medications compared to patients with low risk. Since there is no specific risk stratification for patients with STEMI, we suggested physicians might have different perceptions on the

benefit-risk of using guideline-recommended medications for different patient groups, hence a more personalized approach is required.

In **Chapter 5**, a qualitative study was conducted to elicit physician's perspective on factors influencing the decision to prescribe statins as part of CVD prevention and control. These factors were mapped into a research-based framework analysing the use of statins by general practitioners in England. We found factors operating at microlevel (patient's characteristics, physician's professional experience) and macrolevel (NHIS, international references) contributed to physicians' decision making to prescribe statins. We could not identify any factors from mesolevel as observed in England. It was concluded that the role of primary healthcare in CVD prevention was still limited, because of the relatively recent introduction of the NHIS in Indonesia.

In **Chapter 6**, disparities in the effects of statins on lipid parameters between Dutch men and women were investigated in an inception cohort study using the PharmLines Initiative database. Limited by the small sample size, we found that regardless of the presence of previous CVD, statin therapy increased the level of high-density lipoprotein cholesterol (HDL-c) from baseline more significantly in women than in men. However, the proportion of both men and women who achieved the guideline-recommended low-density lipoprotein cholesterol (LDL-c) target was less than 40% suggesting more investigation on the interplay between statin dose and the level of adherence in statin users that might influence the achievement of this target.

In **Chapter 7**, an inception cohort study using the same database, the association between adherence to statin therapy and lipid parameter response was estimated in first-time statin users on standard dose and low dose. We found the level of adherence to statin therapy was similarly associated with the LDL-c response between standard- and low-dose users. However, with regards to sex, the same level of adherence was associated with a significantly faster rate of reduction of LDL-c in women than in men. This disparity between sexes was also observed in the subgroup of statin standard-dose, but not in the subgroup of low-dose.

In **Chapter 8**, we described how our findings may influence clinical practice and future research. It is apparent that more investigations in the form of longitudinal and quantitative studies are needed to estimate the burden of CVD and the use of

preventive medications in Indonesia. The local clinical guideline needs to be more promoted and its implementation needs to be optimized by addressing factors hindering the use of guideline-based preventive CV medications, including the needs of risk stratification to guide the treatment. Opportunities to enhance the role of primary healthcare in CVD prevention and control in Indonesia within the new NHIS scheme needs to be discussed and established. In the Dutch setting, there is a need for a slight improvement on the achievement of the target using statins in order to optimize CVD prevention. A database linkage between population-based and dispensing database provides an opportunity to investigate more important variables (e.g. patients' adherence, dose of medications) influencing the optimal use of CV preventive medications such as statins in the Netherlands. The possibility of sex disparities in the effect of statins on lipid parameters needs further study with a larger sample size including patients' level of adherence and the fluctuation of statin dose over the period of follow-up.

Samenvatting

Hart- en vaatziekten (Engels: CardioVascular Diseases (CVD)) behoren tot de hoofdoorzaak van ziekte en sterfte wereldwijd. Meer dan 75% van CVD-gerelateerde sterfgevallen, met name door coronaire hartziekten (CHD) en beroertes, vinden plaats in lage- en middeninkomenslanden (LMICs), waaronder Indonesië. Hoewel CVD niet langer de hoofdoorzaak is van ziekenhuisopnames en sterfgevallen in Nederland, een van de hoge-inkomenslanden (HILs), is het nog steeds een hoofdoorzaak van sterfgevallen in Indonesië. Om de druk van CVD te verlichten, hebben de World Health Organization (WHO) en internationale cardiovasculaire (CV-)gerelateerde organisaties klinische richtlijnen opgesteld die onder andere aanraden om preventieve CV-medicijnen te gebruiken gebaseerd op een individuele CV risicobeoordeling. Er zijn verschillende klassen van preventieve CV-medicijnen die aanbevolen worden met bewezen effectiviteit in het voorkomen van CVD. Deze medicijnen zijn antitrombose-middelen (aspirine, P₁₂Y₁₂ inhibitors, antistollingsmiddelen), antihypertensie-middelen (bijv., bètablokkers, remmers van angiotensineconverterend enzym [ACE-remmers], angiotensine receptor blokkers [ARBs]) en cholesterolverlagende medicamenten, zoals statines. Het blijft een uitdaging om het gebruik van deze preventieve medicijnen te optimaliseren in een risicopopulatie. Deze uitdagingen zijn anders in Indonesië als LMIC dan in Nederland als HIL. Onderzoek naar de effectiviteit van deze medicijnen in de CVD-risicopopulatie van Indonesië is erg beperkt. De primaire gezondheidszorgsector in de nieuwe National Health Insurance System (NHIS, namelijk de *Jaminan Kesehatan Nasional-Kartu Indonesia Sehat* [JKN-KIS]) in Indonesië, speelt een belangrijke, invloedrijke rol in het gebruik van preventieve CV-medicijnen. In Nederland is systematische en programmatische CVD-preventie echter al in 2005 geïntroduceerd. De Nederlandse primaire gezondheidszorgsector en zorgverzekeringsmaatschappijen spelen een belangrijke rol bij CVD-preventie. De klinische richtlijnen die risicofactoren voor CV toepassen, zijn al in ontwikkeling sinds mid-2000. Het verschil tussen de stadia van CVD-preventie in beide landen kan resulteren in specifieke uitdagingen qua gebruik van preventieve CV medicijnen in Indonesië en Nederland. Deze dissertatie beoogt de huidige uitdagingen in het gebruik van preventieve CV-medicijnen op basis van aanbevolen richtlijnen in Indonesië en Nederland te identificeren. CHD is het hoofdtype CVD waar we in ons onderzoek op focussen. Deze dissertatie is opgedeeld in twee gedeeltes: **Deel 1 (Hoofdstuk 2-5)** onderzoekt de langdurige CVD druk bij Aziaten en de uitdagingen bij het gebruik van preventieve CV-medicijnen in Indonesië, en **Deel 2 (Hoofdstuk 6**

en 7) onderzoekt de uitdagingen bij het gebruik van preventieve CV-medicijnen in Nederland.

In **Hoofdstuk 2** is een systematische evaluatie en meta-analyse beschreven om vast te stellen wat het lange-termijnrisico (> 10 jaar) is van CVD en de risicofactoren in Aziatische bevolkingsgroepen. De meeste onderzoeken die hier omschreven worden, zijn toegespitst op de Oost-Aziatische regio, geen enkele kwam uit Indonesië. Het lange-termijnrisico van fatale CVD in de Aziatische bevolkingsgroep is 6.35% tijdens een gemiddelde duur van 20 jaar. Het lange-termijnrisico van een fatale beroerte is hoger dan dat van fatale CHD. Verschillende risicofactoren tussen CHD en beroertes worden omschreven. Risicofactoren voor fatale CVD die geïdentificeerd zijn met behulp van de meta-analyse zijn: het mannelijk geslacht, hogere leeftijd, roken. In tegenstelling tot voorgaande onderzoeken onder de Westerse bevolking, kwamen wij tot de conclusie dat andere methodes misschien nodig zijn voor CVD-preventie en controle in Aziaten vanwege de verschillende uitkomsten en risicofactoren in vergelijking met Westerlingen.

Hoofdstuk 3 beschrijft een cohortonderzoek met als doel de effectiviteit vast te stellen van preventieve CV-medicijnen op basis van aanbevolen richtlijnen ten aanzien van ziekenhuissterfte bij patiënten met hartinfarct met ST-elevatie (STEMI), een acute manifestatie van CHD. Deze studie is uitgevoerd met behulp van het Jakarta Acute Coronary Syndrome (JAC) Register. We zien het gebruik van preventieve medicijnen bij 42% van de patiënten met STEMI die niet behandeld werden met 'acute primaire percutane coronaire interventie (pPCI)' therapie, van wie het merendeel laat in het ziekenhuis arriveerde. Eenderde van deze patiënten kregen geen preventieve medicijnen op basis van aanbevolen richtlijnen bij opname in het ziekenhuis. Nadat er rekening is gehouden met mogelijke versturende factoren in de statistische analyse, werd geschat dat het gebruik van preventieve CV-medicijnen in deze bevolkingsgroep het overlijdensrisico in het ziekenhuis kan verlagen met meer dan 50%.

In **Hoofdstuk 4**, met het gebruik van een cohortonderzoek uit hetzelfde register, evalueren we de predictoren voor suboptimaal gebruik van preventieve CV-medicijnen op basis van aanbevolen richtlijnen bij patiënten met STEMI. We ontdekten de volgende predictoren: de aanwezigheid van non-anterior MI; leeftijd > 65 jaar; geen behandeling met acute reperfusetherapie; een familiegeschiedenis met CAD;

een trombolysie in myocard infarct (TIMI) score ≥ 4 . Het niet behandeld zijn met reperfusie was geïdentificeerd als de predictor voor alle geneesmiddelklassen.

In zowel **Hoofdstuk 4** als **5** leek een treatment-risk paradox te bestaan, dit is een situatie waarbij patiënten met een hoger risico minder medicatie krijgen in vergelijking met patiënten met een lager risico. Omdat er geen specifieke risicostratificatie voor patiënten met STEMI is, suggereren wij dat artsen mogelijk verschillende perspectieven hebben op het risico of profijt dat het gebruik van medicijnen op basis van richtlijnen met zich meebrengt voor verschillende patiëntengroepen. Hierdoor is een gepersonaliseerde aanpak nodig.

Het kwalitatief onderzoek in **Hoofdstuk 5** toont het perspectief van een arts op de factoren die van invloed zijn op de beslissing om statines voor te schrijven bij CVD-preventie en controle. Deze factoren zijn gebaseerd op een research-based raamwerk dat het gebruik van statines bij huisartsen in Engeland analyseert. We ontdekten factoren die op microlevel opereren (eigenschappen van patiënten, professionele ervaring van de behandelend arts) en op macrolevel (NHIS, internationale richtlijnen) welke allemaal bijdroegen aan de beslissing van de arts om statines voor te schrijven. We hebben geen factoren kunnen identificeren op mesolevel zoals in Engeland. Er werd geconcludeerd dat de rol van de primaire gezondheidszorg in CVD-preventie nog steeds gelimiteerd is, vanwege de relatief recente invoering van de NHIS in Indonesië.

In **Hoofdstuk 6** worden de verschillende effecten van statines op lipide-parameters tussen Nederlandse mannen en vrouwen onderzocht in een inceptie-cohort studie met gebruik van de PharmLines Initiative database. Ondanks de gelimiteerde grootte van onze testgroep, hebben we toch aan kunnen tonen dat ongeacht de aanwezigheid van eerdere CVD, statine therapie de concentratie van high-density lipoproteïne cholesterol (HDL-c) sinds start van medicatie significanter verhoogde in vrouwen dan in mannen. Echter, het aantal mannen en vrouwen dat de aanbevolen low-density lipoproteïne cholesterol (LDL-c) doel behaalde was voor beide geslachten lager dan 40%. Dit suggereert dat meer onderzoek naar de interactie tussen statine dosering en mate van adherence in statinegebruikers nodig is in het behalen van de doelstelling. Een inceptie-cohortonderzoek met dezelfde PharmLines database is gebruikt in **Hoofdstuk 7** om het verband tussen adherentie aan statinetherapie en lipide parameter response te berekenen bij nieuwe gebruikers voor een standaard dosering

en lage dosering. We ontdekten dat het adherentie niveau voor statine therapie evenwijdig gerelateerd was aan de LDL-c response tussen de standaard- en lage-dosering-gebruikers. Maar als we naar geslacht kijken, zagen we hetzelfde adherentie niveau geassocieerd met een significant snellere reductie van LDL-c bij vrouwen dan bij mannen. Deze ongelijkheid tussen de geslachten werd ook geobserveerd in de subgroep van standaard statine dosering, maar niet in de subgroep van de lage dosering.

In **Hoofdstuk 8** beschrijven we hoe onze bevindingen mogelijk van invloed kunnen zijn op de klinische praktijk en toekomstig onderzoek. Het is duidelijk dat er meer onderzoek moet worden gedaan in de vorm van longitudinaal en kwantitatief onderzoek om de ziekte en sterftelast van CVD en het gebruik en effecten van preventieve medicijnen in Indonesië goed in te kunnen schatten. De lokale, klinische richtlijnen moeten gepromoot worden en de implementatie hiervan moet geoptimaliseerd worden door factoren aan te pakken die het gebruik van preventieve CV-medicijnen op basis van richtlijnen hinderen, waaronder de wens om risicostratificatie te gebruiken bij een behandeling. Mogelijkheden om de rol van de primaire gezondheidszorg te vergroten bij CVD -preventie en bestrijding in Indonesië binnen het nieuwe NHIS systeem, moeten worden besproken en vastgelegd. In Nederland is er behoefte aan verbetering van het halen van de doelstellingen ten aanzien van cholesterol verlaging met statine om CVD-preventie te optimaliseren. De huidige database link tussen Lifelines en de geneesmiddeldatabase van de Universiteit Groningen biedt een mogelijkheid om de belangrijkste variabelen te onderzoeken (bijvoorbeeld adherentie van patiënten, dosering van medicatie) die het optimale gebruik van CV preventie medicijnen beïnvloeden in Nederland. Mogelijke verschillen door geslacht bij het effect van statines op lipide parameters verdient verder onderzoek met een grotere testgroep waarbij ook het adherentie-niveau van patiënten en de fluctuatie van statine dosering tijdens de follow up worden betrokken.

Acknowledgement

I am aware that the PhD track provides interesting challenges and I would not have completed this amazing process without many people who have come into my life and walked by my side. Therefore, I would like to express my gratitude to all these people and institutions who have enabled me to overcome the challenges with the greatest enjoyment below:

My promotores, **Prof. Eelko Hak** and **Prof. Katja Taxis**.

Dear **Eelko**, thank you for giving me many opportunities to start my PhD studies and for your detailed supervisions of my work. I remember the first time I contacted you by email, it was in 2012. We only met once via skype that year. I could not manage the workload at the university where I worked at that time so I could not continue our contact. Then I contacted you again in 2015. It was beyond my expectation that you still remembered me and gave me another chance to continue my halted PhD plan. Even more, you also introduced me to the LPDP scholarship scheme which I was oblivious with. I am really thankful for this introduction. You have taught me about how to write an article in an effective and persuasive way and to critically think about what really matters regarding a research subject and associated methodology. Thank you for improving my manuscript and for connecting me to so many people who later would help me finishing the project. Thank you for your fairness and for always being available to provide prompt responses and imminent support. It was such a great pleasure to work with you and I hope I will have another opportunity to work with you in the future.

Dear **Katja**, thank you for supervising me during my PhD program. Thank you for your warm approach. I learned a lot from your expertise in cardiovascular research in developing countries. You taught me about how to structure my writing, how to present the same idea in more concise and better words. Thank you for helping me to improve the manuscript and for providing a lot of insights on the qualitative study. I am really happy you supervised my work and I do really hope for another opportunity to work with you in the future.

My former second supervisor, **Prof. Bob Wilffert**,

Dear **Bob**, thank you for your warmth greetings in my mother language. I really appreciate it! Thank you for your time to discuss the systematic review during my earlier time as a PhD student. I appreciate your inputs on my writings and prompt response whenever I had difficulties during my PhD time.

The Indonesia Endowment Fund for Education (Lembaga Pengelola Dana Keuangan, LPDP) who sponsored my PhD program.

The reading committee, **Prof. Aukje Mantel-Teeuwisse**, **Prof. Petra Denig** and **Prof. Eric van Roon**, who spent time to review and gave constructive advice for my thesis.

Prof. Geny Groothuis and Tim Zwaagstra, thank you for stopping by at the Universitas Surabaya in 2012, although it was not anywhere in your plan. This way I had an opportunity to be interviewed as a PhD candidate. I don't think I would be here if you did not come by that day!

Prof. Herman J. Woerdenbag, thank you for welcoming my email and referring me to Prof. Eelko Hak.

All my collaborators: **Rizwandy Wasir**, **Dr. Floriaan Schmidt**, **Dr. Atiquil Islam**, **Prof. Erik Buskens**, **Dr. Talitha Feenstra**, **Prof. Maarten Postma**, **Dr. Surya Dharma (Jakarta Acute Coronary Syndrome [JAC] Registry)**, **Nunung Nursyarofah**, **Thang Nguyen**, **Sari Prayudeni**, **Riris Rachmawati**, **Prof. I Wayan Wita**, **Nicholas Hunt**, **Joanna Emmens**, **Jens Bos**, **Prof. Rudolf de Boer**, and **Dr. Stijn de Vos**.

Thank you for all your time and thoughts, for the scientific input on my PhD project and your prompt response resulting in a project that could be finished in time.

Other current and former member of PTE2 and PEGET: **Prof. Maarten Postma**, **Prof. Van Puijenbroek**, **Nynke**, **Thea**, **Pepijn**, **Didik**, **Marcy**, **Neily**, **Jurjen**, **Pieter**, **Ury**, **Qi**, **Tia**, **Lusi**, **Aizati**, **Doti**, **Lan**, **Thang**, **Wandy**, **Eva**, **Christiaan**, **Heleen**, **Linda**, **Khairul**, **Afifah**, **Sofa**, **Fajri**, **Ira**, **Abraham**, **Hans**, **Tanja**, **Indrè**, **Simon**, **Taichi**, **Monik**, **Jens**, and **Felicia**.

Thank you for your support and help during my study.

I also thank **Jannie, Anja, Jugo, Bert and Yvonne** for your warm and technical support in administration and IT department.

My former Dean, **Dr. Christina Avanti**, thank you for inviting Tim and Prof. Gennie at the Universitas Surabaya in 2012 and for informing me about their coming. Otherwise I would not have had this opportunity. Thank you for allowing me to pursue my PhD and for visiting me one time at the University of Groningen and for warm meeting with other colleagues and friends. I really appreciate the moment!

My current dean, **Dr. Farida Suhud**, and my former teacher, **Ardy Winoto** (late), thank you for your warm support during my studies. Thank you for encouragement through your texts and messages. I really appreciate it. You have given me an example on how to live wisely as I am about getting older.

The Founder and Director of the CMIPC/PIOLK Universitas Surabaya, **Dr. Adji Prayitno**, thank you so much for allowing me to pursue my PhD, Pak. Without your approval, I would not be here. Thank you for giving me advise before I went to the Netherlands. I remember that you said everyone has a different way to gain success, it does not depend on whether you are introvert or extrovert (and I am an introvert). Thank you for increasing my confidence. I hope the skills I gained during my PhD can be more helpful for our unit in the future.

My **students, friends, and relatives** who said goodbye at the airport and prayed for me: thank you very much. I remember you all and your support in my heart. I could never make it without you.

My friends during the research in Jakarta: **Daisy, Hadi** and **Yovita**, thank you for helping and accompanying me in difficult times. Your time and sincere support is much appreciated!

My friends in Surabaya and other places in Indonesia: **Lisa, Christine, Yenny, Fang Chu, Yola, Yemima**, thank you for always praying for me. **Linda** and **Lantin**, thank you for your generous time and ears!

My spiritual mentor in Indonesia and in Groningen: **Kak Iis, Kak Harry, Kak**

Getha, Richard, Ria, Rosbina, thank you for providing such a great support through time, messages, listening ears, and prayers. Richard, Ria, and Rosbina, thank you for inviting me at your place in Groningen. I would never forget such experiences and I hope I can visit you again someday in the future!

My officemates, **Ivan, Akbar** and **Yuan**. Thank you for being partners in crime in the happy and frustrating times during my PhD. You have provided a very encouraging environment to learn, to debate, to share, to argue, to build great moments together. It was really a great pleasure to get to know you all and to work with you. I hope in the future we have an opportunity to work together and to chat like in the old days.

My paranympths, **Sofa** and **Yuan**, thank you for being my paranympths. Thank you for sharing your time and stories with me. I wish you and your family a good and successful life!

My friends in Groningen:

Marcy, Ira, Cecilia Hennep, Alice and **Jinna**, thank you for sharing every moment with me and listening to my venting. Thank you for your prayer and support during my PhD. I am grateful to know you and to learn a lot of things from you all.

Mark and **Snowy**, thank you for every moment. I really appreciate it. You have taught me about work-life balance, love and life in the Netherlands. It was beautiful and one of the bests!

My sisters and brothers at CMIPC/PIOLK, Indonesia:

Yosi, Cecilia, Fauna, Ika and **Steven** thank you for your support and listening ears during my PhD. Thank you for giving me insights and encouragement on how to manage my PhD project in more effective way.

Bobby, thank you for your friendship, support and advise on life and my PhD journey, for our trips across the Netherlands. I really appreciate the moments! I wish you a great success with your PhD journey!

Ming Ming, thank you for your encouragement to start pursuing a PhD. Thank you for introducing me to the term 'Pharmacoepidemiology' and to refer me to Prof. Herman. You have inspired me to work hard during my PhD. I really appreciate your

visit to Groningen and your never ending support! I wish you a great success with your PhD journey!

My family:

Mama and Papa, I am grateful and proud to have you both as my parents. I would not exist without your love. I would not be who I am now without your sacrificial love. The word 'thank you' will not be enough. I wish you a good and joyful life. To my brother, **Rizal**, thank you for your prayer and support. May we all create a loving and meaningful moment together at every given opportunity.

Finally, I thank God that all this happened and that I was given an opportunity to meet, to know, and to live with you all for a moment in this world. I am really grateful to know you all. May you all have a blessed and successful life in the year to come.

ABOUT THE AUTHOR

Sylvi Irawati was born in Malang, East Java, Indonesia. She obtained her bachelor's degree in Pharmacy at the Faculty of Pharmacy, Universitas Surabaya, Surabaya, Indonesia in 2003. She finished her professional degree in Pharmacy, major in Clinical Pharmacy, in 2004 at the same university. She worked as a researcher at the Department of Research & Development in a local pharmaceutical industry in Surabaya, formulating liquid vitamins and injections, from 2004 to 2006. She, then, pursued her master's degree in Clinical Pharmacy at the same university from 2006 to 2008. She graduated with *cum-laude* from all her educations. She has been working as a drug information pharmacist and researcher at the Centre for Medicines Information & Pharmaceutical Care (CMIPC/Pusat Informasi Obat dan Layanan Kefarmasian, PIOLK), Universitas Surabaya, since 2007 and as a lecturer at the Department of Clinical & Community Pharmacy, Faculty of Pharmacy of the university since 2011. She is a member of the Committee of Pharmacy and Therapeutic and a visiting clinical pharmacist at the St. Vincentius a Paulo Hospital, Surabaya, since 2009, through a collaboration between CMIPC and the hospital. Her interest in research and lecture is mainly in the use of cardiovascular medications in the hospital and community. She started her PhD track under the supervision of Prof. Eelko Hak and Prof. Katja Taxis in March 2016 at the Department of Pharmacotherapy, -Epidemiology & -Economics (PTE2), University of Groningen, the Netherlands. Her PhD project was funded by the Indonesia Endowment Fund for Education (Lembaga Pengelola Dana Pendidikan, LPDP) and focused on the challenge in the use of cardiovascular medications in Indonesia and the Netherlands. After completing her PhD, she will continue her work at Universitas Surabaya.

PhD PORTFOLIO

Name	:	Sylvi Irawati
Unit of PhD	:	Department of Pharmaco-Therapy, -Epidemiology & -Economics, Groningen Research Institute of Pharmacy (GRIP), University of Groningen, the Netherlands.
PhD period	:	2016-2020
Promotors	:	Prof. Eelko Hak Prof. Katja Taxis

Courses/ Workshops	Year	Workload (hours/ ECTS)
Internal		
Data handling and pharmacoepidemiology in practice	2016	5
Pharmacoeconomics	2016	5
Presentation skills	2016	0.5
Systematic review and meta-analysis	2016	1
Scientific integrity	2016	2.5
Pharmacoepidemiology UK	2016	5
Managing your PhD	2016	2
Basic Medical statistics	2017	3
Study design in clinical epidemiology	2017	4
Science writing	2017	2
Publishing in English	2017	3
Advance topics in pharmacoepidemiology	2017	5
Epidemiology and applied statistics	2018	3
Quality-of-Life and Patient-Reported Outcome Measures (QoL & PROMs)	2019	2.5

External

Medical Sciences Summer School, University Medical Center Groningen, Groningen, the Netherlands: fundamentals of biobanking and cohort research.	2017	Not available
Pre-conference courses of the 34th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ICPE), Prague, Czech Republic: Advanced drug utilization research Using field studies to value-add in pharmacoepidemiology Best practices for designing, implementing and evaluating risk minimization programs	2018	Not available
European Consortium for Political Research (ECPR), Budapest, Hungary: survey and questionnaire design.	2019	4

Published and submitted article in scientific journals

Irawati S, Wasir R, Schmidt AF, Islam A, Feenstra T, Buskens E, Wilffert B, Hak E. Long-term incidence and risk factors of cardiovascular events in Asian populations: systematic review and meta-analysis of population-based cohort studies. *Curr Med Res & Opin* 2019; 35:2, 291-9.

Irawati S, Dharma S, Taxis K, Thang N, Nursyarofah N, Wilffert B, Hak E. Association between adherence to guideline-recommended preventive medications and in-hospital mortality among non-reperfused ST-elevation myocardial infarction patients admitted to a tertiary care academic center in a developing country. *Glob Heart* 2020; 15(1): 8.

Irawati S, Prayudeni S, Rachmawati R, Wita IW, Wilffert B, Hak E, Taxis K. Key factors influencing the prescribing of statins: a qualitative study among physicians working in primary healthcare facilities in Indonesia. *BMJ Open* 2020; 10(6): e035098.

Irawati S, Dharma S, Taxis K, Nguyen T, Nursyarofah N, Wilffert B, Hak E. Predictors for prescribing guideline-recommended medications at-discharge in patients with acute ST-segment elevation myocardial infarction (STEMI) admitted to a tertiary care academic center in a developing country. (*submitted*)

Irawati S, Emmens JE, de Vos S, Bos JHJ, de Boer R, Hak E. Association between adherence to statin therapy and low-density lipoprotein cholesterol (LDL-c) response in first-time users of standard-dose and low-dose statins: the PharmLines Initiative. (*submitted*)

Hunt NB, Emmens JE, **Irawati S**, Bos JHJ, Wilffert B, Hak E, de Boer R. Sex disparities in the effect of statins on lipid parameters: the PharmLines Initiative. (*submitted*)

Wasir R, **Irawati S**, Makady A, Postma M, Goettsch W, Buskens E, et al. 2019. Use of medicine pricing and reimbursement policies for universal health coverage in Indonesia. *PLoS ONE* 2019; 14(2): e0212328.

Wasir R, **Irawati S**, Makady A, Postma M, Goettsch W, Feenstra T, et al. 2019. The implementation of HTA in medicine pricing and reimbursement policies in Indonesia: Insights from multiple stakeholders. *PLoS ONE* 2019; 14(11): e0225626.

Mulyono I, **Irawati S**, Susilo AP, Claramita M. Pharmacist-patient communication in Indonesia: the Roter Interaction Analysis System (RIAS) in socio-hierarchical context. *Pharm Edu* 2019(1): 359-69.

